

Original Article

A Comparative Evaluation of the Antioxidant and Phytochemical Properties of Various Extracts of *Ocimum Sanctum* Linn Grown in Jaffna District, Sri Lanka

Gowri Rajkumar^{1*}, Vilochana Alahendra¹, Vinotha Sanmugarajah²

¹Department of Botany, Faculty of Science, University of Jaffna, Sri Lanka

²Unit of Siddha Medicine, University of Jaffna, Sri Lanka.

Received: 16.11.2024; Revised: 18.12.2025; Accepted: 28.12.2025; Published: 15.01.2025

Abstract

Background and Aim: *Ocimum sanctum* Linn (*O. sanctum*) (Lamiaceae) is a valuable medicinal herb. It contains several therapeutic characteristics due to its bioactive components. This study aimed to compare the phytochemical and antioxidant properties of different extracts of the leaves and flowers of the *O. sanctum* plant, which grows in Sri Lanka.

Materials and Methods: Standard procedures were followed for the qualitative and quantitative phytochemical analyses and the DPPH as well as ABTS antioxidant activities.

Results: The findings demonstrated that the ethanolic and methanolic extracts had adequate concentrations of flavonoids, phenols, tannins, and alkaloids compared with the aqueous extracts of the flowers and leaves of *O. sanctum*. The supreme quantity of the flavonoid content existed in the ethanolic flower extracts (50.88 ± 0.62 μg QE/g) and methanolic leaf extract (49.11 ± 0.27 μg QE/g), while the higher amount of the phenol content (20.96 ± 0.65 μg GAE/g) was found in the ethanolic leaf extract. The methanolic leaf extract had the highest tannin content (409.95 ± 0.66 μg TAE/g) compared with other extracts. Moreover, the highest amount of alkaloid was observed in flowers (75.83 ± 0.55 mg/g) and not in leaves (52.60 ± 0.66 mg/g). Furthermore, the methanolic flower extract exhibited the greatest DPPH (IC_{50} : 0.68 ± 0.19 $\mu\text{g}/\text{mL}$) and ABTS (IC_{50} : 1.82 ± 0.32 $\mu\text{g}/\text{mL}$) activities compared with the Trolox.

Conclusion: The present study proved that although the leaf extracts contained higher concentrations of phytochemicals, the methanolic flower extract of *O. sanctum* had a greater potential for antioxidants than the methanolic leaf extract.

Keywords: Antioxidant, Comparative evaluation, Various extracts, *Ocimum sanctum*, Phytochemicals, Sri Lanka.

***Corresponding Author:** Gowri Rajkumar, Department of Botany, Faculty of Science, University of Jaffna, 40 000, Sri Lanka, Email: gowrir@univ.jfn.ac.lk.

©2025 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited.

Introduction

Therapeutic benefits of various diseases might be provided by chemical structures and bioactive composites (therapeutic ingredients) present in medicinal plants (1). Approximately, 80% of people worldwide still use traditional medicines such as herbal remedies (2, 3). Herbal medicine is an efficient and safe method of treating diseases and infections. Many people worldwide, particularly those living in developing countries, continue getting their medical care from herbal remedies. Even though synthetic drugs are widely available and effective in treating a wide range of disorders in today's culture, some individuals choose to employ traditional folk remedies since they have less adverse effects (4). Nearly 13,000 secondary metabolites, or phytochemicals, including alkaloids, flavonoids, tannins, and phenolic compounds, have been found in medicinal plants. Secondary metabolites in plants act as defensive chemicals or fulfill specific roles, and have healing benefits (pharmacological actions), such as anti-diabetic, analgesic and antioxidant actions (5-8). These natural phytochemicals of many therapeutic plants can help reduce inflammation, protect against oxidative damage, and improve liver function. Moreover, they are mostly determined by the place in which they are grown, the environment of that region, and the technique as well as season of collection (9).

Mahmood (2012) states that *Ocimum sanctum* Linn (*O. sanctum*), often known as holy basil or *tulsi*, is an excellent exemplar of nature's pharmacy (10). Tulsi has received a lot of attention in traditional medicine for its rich phytochemical composition and various pharmacological actions in treating a variety of health problems. This perennial plant is highly appreciated in traditional medical systems for its antioxidant, antibacterial, anti-inflammatory, anti-asthmatic, immune-modulatory, anti-stress characteristics and anti-carcinogenic effects (10, 11). Its distinct parts have long been utilized in traditional medicine to manage the conditions, such as bronchitis, malaria, diarrhea, skin conditions, and rheumatism (12). *O. sanctum* has been linked to a number of therapeutic properties (13), and is commonly found in various Asian countries, including India, Sri Lanka, Himalaya, etc. (14, 15). Various chemical components resembling chavicol and linalool present in essential oil derived from *O. sanctum* leaves and flowers are efficiently used in herbal medicine (16).

O. sanctum is commonly utilized in herbal medications in management of various ailments in the Sri Lankan Traditional System of Medicine. It includes exceptional elements with great biological activities. Nonetheless, more scientific research is required to establish additional therapeutic potentials and pharmacological effects. No research has been published in Sri Lanka, on the efficiency this plant's phytochemicals and antioxidants utilizing comparable laboratory and scientific methodologies to date. Thus, the purpose of this study was to evaluate the phytochemical and antioxidant potentials of several leaf and flower extracts from the *O. sanctum* plant, which grows in Sri Lanka.



Figure 1. Leaves and Flowers of the *Ocimum Sanctum* L. Plant.

Materials and Methods

Collection of Samples

The *O. sanctum* plant's leaves and flowers were collected in December 2022 from the Jaffna District in Sri Lanka's Northern Province. The created herbarium underwent botanical authentication, and then the voucher specimen was placed in the Department of Botany, Faculty of Science, University of Jaffna.

Preparation of the Samples

To prevent the direct loss of phytoconstituents from sunshine, the samples were carefully allowed to air-dry at room temperature for three weeks after being frequently cleaned with tap water to remove dust and grime (Figure 1-3). The



Figure 2. The Plant Herbarium of *Ocimum sanctum* L.

Subsequently, they were filtered through Whatman No.1 filter paper. The part of the filtered content was concentrated using rotatory evaporator (Buchi), and the other part was kept in refrigerator at 4°C for further uses. The analysis was done for three replicates of each extract for both samples.

Qualitative and Quantitative Phytochemical Analysis



Figure 3. Powder Samples of the Leaves and Flowers of the *Ocimum Sanctum*.

materials that had been shade-dried were ground into a fine powder using a pulverizer and sieved through as many as 80 meshes. For further examination, it was stored separately in sealed bottles at 31±3°C.

Extraction Procedure

Each sample was extracted with ethanol, methanol and water separately using the cold extraction technique (17). A total of 50 g of powdered materials of each sample was separately weighed and placed in 500 ml of culture bottles. 150 ml of 100 % absolute ethanol/ methanol/ water (1:3) was added to it and mixed well separately. Lids of the bottles were covered with parafilm. The solution was kept for 5 days with occasional shaking using shaker at 150 rpm for 15 minutes in every morning and evening.

The ethanol, methanol, and aqueous extracts of every sample of *O. sanctum* were individually screened for phytochemicals such as flavonoids, phenols, tannins, alkaloids, terpenoids, saponins, glycosides, anthraquinones, etc., using the standard laboratory techniques mentioned by Rajkumar *et al.* 2021, 2022, and 2023 (17-19).

For quantification purposes, total phenol (µg GAE/g) and tannin (µg TAE/g) contents were estimated based on the Folin-Ciocalteu colorimetric method, (20, 21) total flavonoid content (µg QE/g) based on the aluminum colorimetric method, (22) and total alkaloid contents (mg/g) based on Edeoga *et al.* (2005) (23) and Aliyu *et al.* (2008) (24) in each filtered sample of *O. sanctum* using a spectrophotometer (17).

Evaluation of the Antioxidant Activity

The DPPH (1,1-diphenyl-2-picryl-hydrazyl) and ABTS (2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid)) complementing tests were used to measure the in vitro antioxidant activity of each filtered sample of *O. sanctum* using a spectrophotometer. Trolox served as the standard for comparison.

Utilizing the graph and the standard equation, the free radical scavenging activity (%) and IC₅₀ value were determined.

DPPH Free Radical Scavenging Assay:

The method defined by Yamaguchi *et al.* 1998 (25) with a few adjustments has been used for this determination (26). A series of different solvent extracts of plant parts (100 µl, 200 µl, 300 µl, 400 µl, 500 µl) were prepared separately. 1 ml of DPPH (0.002g/50 ml) in methanol was added in each sample, and the absorbance was measured at 517nm after incubation for 15 min.

$$\text{Inhibition \% radical DPPH activity} = \frac{\text{Absorbance of control} - \text{Absorbance of the sample}}{\text{Absorbance of control}} \times 100$$

ABTS Free Radical Scavenging Assay:

The procedure outlined by Re *et al.* (1999) (27) was used to measure this assay. A mixture of ABTS (7 mM) and potassium persulfate (2.5 mM) was prepared and kept at room temperature for 16 hours in a dark room to produce ABTS⁺. Different solvents of the extracts were mixed with diluted ABTS solution, and the absorbance was measured at 734 nm up to 6 min.

$$\text{Inhibition \% radical ABTS activity} = \frac{(\text{Absorbance of ABTS + methano mixture}) - \text{Absorbance of the sample}}{\text{Absorbance of the sample}} \times 100$$

Statistical Analysis

ANOVA and Tukey's multiple comparisons at probability value ($p \leq 0.05$) were used in the data analysis, which was carried out using Minitab 17 software. Every analysis was carried out in triplicate, and the mean \pm standard deviation (SD) was reported as the result of the analysis.

Results and Discussion

Plants are highly significant basis of potentially bioactive elements for the advance of new chemotherapeutic drugs. In the first stage to achieve this intention, leaves and flowers of *O. sanctum* were subjected to systematic qualitative and quantitative analyses by different extracts to determine the phytochemicals and their total contents.

Identification of the Phytochemicals

The *O. sanctum* was subjected to a preliminary phytochemical screening in order to identify its phytochemicals from its various extracts of the leaves and flowers. Table 1 shows that there were more phytochemicals in the methanolic and ethanolic extracts of *O. sanctum* than in the aqueous extract which might be due to the fact that some chemicals may not be properly soluble in the aqueous solvent. Notably, the leaves contained a more diverse range of phytochemicals than the flowers. Different extracts of *O. sanctum* ensure the richness of phenols, flavonoids, tannins, alkaloids, etc. Anthraquinones were consistently absent in all extracts of both leaves and flowers. These constituents might be possibly

Table 1. Identification of the Phytochemicals in the *O. sanctum*.

Phytochemical	Leaves			Flowers		
	Ethanol	Methanol	Aqueous	Ethanol	Methanol	Aqueous
Flavonoids	++	+++	++	+++	+	+
Phenols	+++	++	-	++	+	-
Tannins	++	+++	+	++	+	-
Alkaloids	++	+	+	+++	+	+
Terpenoids	+	+	+	++	+	++
Saponins	+	-	+	+	-	+
Glycosides	+	+	+	+	+	+
Steroids	+	+	+	+	+	+
Anthraquinones	-	-	-	-	-	-
Xanthoproteins	+	+	+	+	-	-
Quinones	+	-	+	-	-	+
Coumarins	-	-	-	+	-	-
Carboxylic acids	+	-	+	-	-	-
Reducing sugars	-	-	+	+	+	+
Proteins and amino acids	++	+	-	++	++	+

accountable for the biological activities of *O. sanctum*.

The leaves of *O. sanctum* have been a subject of numerous scientific investigations due to their therapeutic potential. The present research is almost comparable to several studies which have focused on the phytochemical composition of the leaves, revealing the occurrence of various secondary metabolites (28-36). The flowers contain essential oils, which contribute to their fragrance and therapeutic properties as antioxidant, and antimicrobial agents (37-39).

Determination of Phytochemicals

Table 2 compares the quantities of four distinct phytochemicals—phenol, flavonoid, tannin, and alkaloid—in the ethanolic, methanolic, and aqueous extracts of *O. sanctum*'s leaf and flower parts. The total content of phytochemicals was attained by using the standard curve for each of these phytochemicals (Figures 4, 5 and 6).

Table 2. Determination of Phytochemicals in the *O. sanctum*.

Samples	Extracts/ Sample	Phenols ($\mu\text{g GAE/g}$)	Flavonoids ($\mu\text{g QE/g}$)	Tannins ($\mu\text{g TAE/g}$)	Alkaloids (mg/g)
Leaves	Ethanollic extract	20.96 ± 0.65	41.19 ± 0.24	286.08 ± 0.37	-
	Methanollic extract	14.89 ± 14.89	49.11 ± 0.27	409.95 ± 0.66	-
	Aqueous extract	8.53 ± 0.48	26.54 ± 0.49	242.12 ± 0.34	-
	Powder sample	-	-	-	52.60 ± 0.66
Flowers	Ethanollic extract	15.53 ± 0.44	50.88 ± 0.62	297.16 ± 0.56	-
	Methanollic extract	7.53 ± 7.53	23.88 ± 0.76	261.00 ± 0.48	-
	Aqueous extract	4.59 ± 0.13	18.05 ± 0.39	140.54 ± 0.61	-
	Powder sample	-	-	-	75.83 ± 0.55

The maximum amount of flavonoid content was present in the ethanolic flower extracts ($50.88 \pm 0.62 \mu\text{g QE/g}$) and methanolic leaf extract ($49.11 \pm 0.27 \mu\text{g QE/g}$), while the higher amount of phenol content ($20.96 \pm 0.65 \mu\text{g GAE/g}$) was found in the ethanolic leaf extract. The methanolic leaf extract showed the highest tannin content ($409.95 \pm 0.66 \mu\text{g TAE/g}$) compared with other extracts. Moreover, the highest amount of alkaloid was present in flowers




Figure 4. Standard Curve for Total Phenolic Content.




Figure 5. Standard Curve for Total Flavonoid Content




Figure 6. Standard Curve for Total Tannin Content

($75.83 \pm 0.55 \text{ mg/g}$) and not in leaves ($52.60 \pm 0.66 \text{ mg/g}$). The present study proved that most of the phytochemicals as phenols, flavonoids and tannins were higher in leaves than flowers of *O. sanctum* plant.

As it has been indicated in Table 3, the results of the present study for total phenol and flavonoid contents for the methanolic leaf extract of *O. sanctum* were not consistent with the results of previous studies conducted by Singh *et al.*, 2014; Pathak & Niraula, 2019; Chaudhary *et al.*, 2020 and Singh *et al.*, 2020 (34, 35, 40, 41). Variances in the preparation process, sample storage

technique, and seasonal or geographic variances in the environment from which plant materials were taken, or the standardization method employed could all be the causes of these discrepancies.

A class of metabolites known as phenolic compounds is produced by plants' secondary pathways. Flavonoids, phenolic acids, tannins, and coumarins are naturally occurring materials that can be found in a variety of plant products (42, 43). According to recent research, phenolic compounds may protect the body against diseases caused by by oxidative stress (42-45).

Table 3. A Summary of Phytochemicals in the Methanolic Leaf Extract of *O. sanctum*.

Phytochemical	The present study	Singh <i>et al.</i> , 2020	Chaudhary <i>et al.</i> , (2020)	Pathak and Niraula, 2019	Singh <i>et al.</i> , 2014
Total phenols ($\mu\text{g GAE/g}$)	14.89 \pm 14.89	703.12 \pm 0.00	87.13 \pm 4.6	180.21 \pm 0.89	4.49 - 9.31
Total flavonoids ($\mu\text{g QE/g}$)	49.11 \pm 0.27	100 \pm 0.0056	221.97 \pm 4.6	67.11 \pm 0.43	14 - 225

Epidemiological studies maintain that extensive period consumption of diets rich in plant polyphenols offer protection against the progress of chronic illness (46-48). Flavonoids are a significant class of natural compounds that are derived from plants and have an extensive array of positive health effects. They are efficiently used as pharmacological, therapeutic and nutraceutical agents. This is explained by their ability to modify the activities of important cellular enzymes as well as their antioxidative, anti-inflammatory, anti-mutagenic, and anti-carcinogenic qualities (49). Plant polyphenols, called tannins, are bitter and astringent agents. They can precipitate or bind to proteins as well as alkaloids (50-52). This tannin-protein complex can provide persistent antioxidant activity (53-55). Alkaloids are predominantly found in plants, with some families of flowering plants having higher concentrations than others (56). Alkaloids generated from plants also have antioxidant properties (57).



In Vitro Antioxidant Activity

Variations in the antioxidant activity of the methanolic extract of leaves and flowers of *O. sanctum* based on DPPH and ABTS assays have been indicated in Table 4 and Figures 7 and 8.

Table 4. DPPH and ABTS Antioxidant Activity of the Methanolic Extract of *O. Sanctum*.

Sample	DPPH ($\text{IC}_{50}\mu\text{g/ml}$)	ABTS ($\text{IC}_{50} \mu\text{g/ml}$)
Leaf	1.95 \pm 0.82	2.59 \pm 0.44
flower	0.68 \pm 0.19	1.82 \pm 0.32

Mean \pm Standard Deviation

**Figure 7.** DPPH Radial Scavenging Activity of Different Parts of *O. Sanctum***Figure 8.** ABTS Radial Scavenging Activity of Different Parts of *O. Sanctum*

Although the leaves and flowers of *O. sanctum* have been documented to possess significant antioxidant potentials, a comparative assessment of their antioxidant activities is essential for understanding their beneficial value.

As stated in Table 4, the maximum radical scavenging activity (as characterized by the lowest IC_{50}) was observed in *O. sanctum* flowers followed by leaves in both DPPH and ABTS assays. Regarding the DPPH test, the methanolic extract of the flowers (IC_{50} : 0.68 \pm 0.188 $\mu\text{g/ml}$) exhibited the best radical scavenging activity in comparison with the methanolic extract of the leaves (IC_{50} : 1.95 \pm 0.82 $\mu\text{g/ml}$) when tested with Trolox. In case of ABTS assay, the flower methanolic extract (IC_{50} : 1.82 \pm 0.32 $\mu\text{g/ml}$) exhibited the best radical scavenging activity in comparison with the leaf methanolic extract (IC_{50} : 2.59 \pm 0.44 $\mu\text{g/ml}$). According to the results, both the leaves and the flowers had significant quantities of antioxidant activity, even though the antioxidant activity of the flowers was higher. Since methanol dissolves many antioxidant compounds efficiently, these assays were performed against methanol extracts. According to the study's findings, the plant's leaves and

flowers might provide a valuable supply of natural antioxidants that could be used to treat degenerative disorders associated with oxidative stress.

Many studies have shown that the *O. sanctum* extract has greater DPPH antioxidant activity than ascorbic acid (58, 59), and that commercially available *O. sanctum* powder demonstrates higher degrees of DPPH radical scavenging activity than vitamin C (60). According to Chaudhary *et al.* (2020) (35), the methanolic leaf extract of *O. sanctum* exhibited antioxidant activity as demonstrated by the DPPH and ABTS assays. The main way that antioxidants work is by scavenging free radicals, which either prevents or delays cellular damage. Natural substances found in dietary supplements and medicinal plants are increasingly being used as therapeutic antioxidants (61). An *in vivo* study indicated that the potential antistressor activity of *O. sanctum* could be partially attributable to its antioxidant properties (62).

Conclusion

The findings of the present study revealed that the ethanolic and methanolic extracts had greater concentrations of flavonoids, phenols, tannins, and alkaloids than the aqueous extracts of the flowers and leaves of *O. sanctum*. The extreme quantity of flavonoid content was found in the ethanolic flower extract and methanolic leaf extract, while the advanced quantity of phenol content was found in the ethanolic leaf extract and the highest tannin content was present in the methanolic leaf extract. Moreover, the highest amount of alkaloid was present in flowers than in leaves, and the methanolic flower and leaf extracts exhibited considerable DPPH and ABTS radical scavenging activities related to the Trolox. This comparative study proved that despite the fact that the leaf extracts contained higher concentrations of phytochemicals, the methanolic flower extract of *O. sanctum* had a better potential for having antioxidant properties than the methanolic leaf extract. Thus, it could be used as a potential therapeutic ingredient in the development of functional medications and its utilization would be a viable alternative to combat various non-communicable diseases.

Acknowledgment

None.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Sofowora A, Ogunbodede E, Onayade A. The role and place of medicinal plants in the strategies for disease prevention. *Afr J Tradit Complement altern med.* 2013; 10(5): 210-229.
- WHO. Research Guidelines for Evaluating the safety and Efficacy of Herbal Medicines, World Health Organization, Manila;1992.
- Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2014; 4:177.
- Yuan H, Ma Q, Ye L, Piao G. The traditional medicine and modern medicine from natural products. *Molecules.* 2016; 21(5): 559.
- Tungmunthum D, Thongboonyou A, Pholboon A, Yangsabai A. Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview. *Medicines.* 2018; 5(3): 93.
- Tran N, Pham B, Le L. Bioactive compounds in anti-diabetic plants: From herbal medicine to modern drug discovery. *Biology.* 2020; 9(9): 252.
- Shehadeh MB, Suaifan GA, Abu-Odeh AM. Plants secondary metabolites as blood glucose-lowering molecules. *Molecules.* 2021; 26(14): 4333.
- Niaz A, Adnan A, Bashir R, Mumtaz MW, Raza SA, Rashid U, Tan CP, Tan TB. The In Vitro α -Glucosidase Inhibition Activity of Various Solvent Fractions of *Tamarix dioica* and 1H-NMR Based Metabolite Identification and Molecular Docking Analysis. *Plants.* 2021;10(6):1128.
- Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants.* 2017; 6(4): 42.
- Mahmood K. Antibacterial activity of essential oil of *Ocimum sanctum* L. *Mycopath.* 2012; 6 (1 & 2): 63-65.
- Singh JP, Singh AK, Bajpai A, Ahmad IZ. Characterization of different *Syzygium cumini* skeys accessions based on physico-chemical attributes and phytochemical investigations. *Int J Pharm Pharm Sci.* 2015; 7(5): 158-164.
- Pattanayak P, Behera P, Das D, Panda SK. *Ocimum sanctum* Linn. A reservoir plant for therapeutic applications: An overview. *Pharm Rev.* 2010; 4(7): 95-105. doi: 10.4103/0973-7847.65323. PMID: 22228948; PMCID: PMC3249909.
- Sen P. Therapeutic potential of Tulsi: From experience to facts. *Drug News Views.* 1993; 1: 15-21.
- Sajjan. Therapeutic Uses of "Wonder Herb" Tulsi (*Ocimum sanctum* Linn) - A Review. *Research Biotica.* 2021; 3(1): 57-62.
- Datiles MJ, Acevedo-Rodríguez P. *Ocimum tenuiflorum* (holy basil). *Cabicompendium.* 110287, CABI Compendium. 2022. doi:10.1079/cabicompendium.110287, CABI International.
- Kalkhorani NM, Dadgar M, Rezaei MB, Hero Abadi F. Comparison of Essential oil of *Ocimum sanctum* L. from Fresh and Dry Aerial Parts by Hydro-distillation and Steam Distillation. *J Med Plan By-products.* 2016; 1: 45-50.

17. Rajkumar G, Jayasinghe MR, Sanmugarajah V. Comparative Analytical Study of Phytochemicals in Selected Antidiabetic Medicinal Plant Seeds in Sri Lanka, *Pharm Sci Res.* 2021; 8(3): 145-155. Available at: <https://scholarhub.ui.ac.id/psr/vol8/iss3/5>
18. Rajkumar G, Panambara PAHR, Sanmugarajah V. Comparative Analysis of Qualitative and Quantitative Phytochemical Evaluation of Selected Leaves of Medicinal Plants in Jaffna, Sri Lanka. *Borneo J Pharm.* 2022; 5(2): 93-103. doi: <https://doi.org/10.33084/bjop.v5i2.3091>
19. Rajkumar G, Sanmugarajah V, Jayasinghe MR. Comparative Phyto and Physicochemical Parameters of the Therapeutic Plant *Syzygium cumini* (L.) Skeels in Jaffna District. *Current Scientia, Faculty of Applied Sciences, University of Sri Jayewardenepura.* 2023; 26 (02): 64-77. <https://doi.org/10.31357/vjs.v26i02.6808>
20. Singleton VL, Orthofer R, Raventos RML. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Meth Enzymol.* 1999; 299: 152-178.
21. Kavitha Chandran CI, Indira G. Quantitative estimation of total phenolic, flavonoids, tannin and chlorophyll content of leaves of *Strobilanthes kunthiana* (Neelakurinji). *J Med Plan.* 2016; 4(6): 282–286.
22. Badarinath AV, Mallikarjuna Rao K, Chetty CMS, Ramkanth S, Rajan TVS, Gnanaprakash K. A Review on in-vitro antioxidant methods: Comparisons, correlations and considerations. *Int J Pharm Tech Res.* 2010; 2(2): 1276-1285.
23. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *Afr J Biotechnol.* 2005; 4(7): 685-688.
24. Aliyu AB, Musa AM, Oshanimi JA, Ibrahim HA, Oyewale AO. Phytochemical analysis and mineral elements composition of some medicinal plants of Northern Nigeria. *Niger J Pharm Sci.* 2008; 7(1): 119– 125.
25. Yamaguchi T, Takamura H, Matoba T, Terao J. HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-2-picrylhydrazyl. *Biosci Biotechnol Biochem.* 1998; 62(6): 1201-4. doi: 10.1271/bbb.62.1201. PMID: 9692204.
26. Blois MS. Antioxidant determination by use of stable free radical. *Nature.* 1958; 181: 1199-1200.
27. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M. Antioxidant activity applying an improved ABTS radical cation decolorization Assay. *Free Radic Biol Med.* 1999; 26: 1231-37.
28. Joshi B, Prasad Sah G, Bahadur Basnet B, Raj Bhatt M, Sharma D, Subedi K, Pandey J, Malla R. Phytochemical extraction and antimicrobial properties of different medicinal plants: *Ocimum sanctum* (Tulsi), *Eugenia caryophyllata* (Clove), *Achyranthes bidentata* (Datiwan) and *Azadirachta indica* (Neem). *J Microbiol Antimicrob.* 2011; 3(1): 1-7.
29. Rana Md. M, Sayeed Md. A, Nasrin Mst. S, Islam M, Rahman Md. M, Alam MF. Free radical scavenging potential and phytochemical analysis of leaf extract from *Ocimum sanctum* Linn. *J Agri Tech.* 2015; 11(7): 1615-1623.
30. Borah R, Biswas SP. Tulsi (*Ocimum sanctum*), excellent source of phytochemicals. *Int J Environ Agric Biotechnol.* 2018; 3 (5).
31. Xia KZ, Perveen N, Khan NH. Phytochemical analysis, antibacterial and antioxidant activity determination of *Ocimum sanctum*. *Pharma Pharmacol Int J.* 2018; 6(6): 490-497. DOI: 10.15406/ppij.2018.06.00223
32. Harichandan SS, Priyadarshini, Kumar SA, Sakshi G, Rahul N. Phytochemical screening and antioxidant activity of methanolic extract of *Ocimum sanctum* Linn. Leaves. *GSC Biol Pharm Sci.* 2019; 08(02): 022-033. DOI: 10.30574/gscbps.2019.8.2.0131
33. Praveen G, Rajesh G. Phytochemical screening and quantitative estimation of total flavonoids of *Ocimum sanctum* in different solvent extract. *Pharma Innov J.* 2019; 8(2): 16-21.
34. Pathak I, Niraula M. Assessment of Total Phenolic, Flavonoid Content and Antioxidant Activity of *Ocimum sanctum* Linn. *J Nepal Chem Soc.* 2019; 40: 30-35.
35. Chaudhary A, Sharma S, Mittal A, Gupta S, Dua A. Phytochemical and antioxidant profiling of *Ocimum sanctum*. *J Food Sci Technol.* 2020; 57(10): 3852-3863. doi: 10.1007/s13197-020-04417-2. Epub 2020 Apr 11. PMID: 32903995; PMCID: PMC7447722.
36. Roy M, Chandnanai Y, Rooplai B, Roy S. Phytochemical Analysis of *Ocimum sanctum* L. Leaf Extracts. *J Ind Vet Assoc.* 2021; 19 (1): 70 – 74.
37. Nasim Milani Kalkhorani, Mohsen Dadgar, Mohammad Bagher Rezaei and Fatemeh HeroAbadi. Comparison of Essential oil of *Ocimum sanctum* L. from Fresh and Dry Aerial Parts by Hydro-distillation and Steam Distillation. *J Med Plan and By-products.* 2016; 1: 45-50.
38. Rakesh Kumar Joshi. Phytoconstituents, traditional, medicinal and bioactive uses of Tulsi (*Ocimum sanctum* Linn.): A review. *J Pharmacog Phytochem.* 2017; 6(2): 261-264.
39. Kumar R, Saha P, Lokare P, Datta K, Selvakumar P, Chourasia A. A Systemic Review of *Ocimum sanctum* (Tulsi): Morphological Characteristics, Phytoconstituents and Therapeutic Applications. *Int J Res App Sci Biotechnol.* 2022; 9(2): 121-126.
40. Singh J, Neha M, Sinha S. Comparison of Total Flavonoid, Phenolic Content and Antioxidant capacity In Leaf and Seed Extracts from White Holy Basil (*Ocimum Sanctum*). *Int J Appl Biol Pharm Technol.* 2014; 5 (4): 34-42.
41. Singh S, Gupta A, Verma S. Phytochemical Screening, TPC, TFC And Antibacterial Activity Against Uti Pathogen of *Ocimum sanctum* Plant Leaf Hydro Alcoholic Extract. *Palarch's J Arch Egyptol.* 2020; 17(7): 13505 – 13513.
42. Lin D, Xiao M, Zhao J, Li Z, Xing B, Li X, Kong M, Li L, Zhang Q, Liu Y, Chen H, Qin W, Wu H, Chen S. An Overview of Plant Phenolic Compounds and Their Importance in Human Nutrition and Management of Type 2 Diabetes. *Molecules.* 2016; 21(10): 1374. doi: 10.3390/molecules21101374. PMID: 27754463; PMCID: PMC6274266.
43. Luna-Guevara Ma. L, Luna-Guevara JJ, Hernández-Carranza P, Espinosa HR, Ochoa-Velasco CE. Phenolic Compounds: A Good Choice Against Chronic Degenerative Diseases. Chapter 3. *Stud Nat Prod Chem.* 2018; 59: 79-108. <https://doi.org/10.1016/B978-0-444-64179-3.00003-7>
44. Cáceres-Jiménez S, Ordóñez-Díaz JL, Moreno-Rojas JM, Pereira-Caro G. 9 – Bioaccessibility and bioavailability of marine polyphenols. *Marine Phenolic Compounds.* Sci Eng. 2023: 265-298. <https://doi.org/10.1016/B978-0-12-823589-8.00008-X>
45. Gadnayak A, Dehury B. Phytochemicals: recent trends in food, pharmacy, and biotechnology, Chapter 5, *Rec Fron Phytochem Appl Food Pharm Cos Biotechnol.* 2023: 85-93. <https://doi.org/10.1016/B978-0-443-19143-5.00031-1>
46. Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev.* 2009; 2(5): 270-8. doi: 10.4161/oxim.2.5.9498. PMID: 20716914; PMCID: PMC2835915.
47. Hano C, Tungmunthum D. Plant Polyphenols, more than Just Simple Natural Antioxidants: Oxidative Stress, Aging and Age-Related Diseases. *Medicines (Basel).* 2020; 9; 7(5): 26. doi: 10.3390/medicines7050026. PMID: 32397520; PMCID: PMC7281114.
48. Natalia VZ, Zubova MY, Nechaeva TL, Kazantseva VV, Goncharuk EA, Katanskaya VM, Baranova EN, Aksenova MA. Polyphenols in Plants: Structure, Biosynthesis, Abiotic Stress Regulation, and Practical Applications (Review). *Int J Mol Sci.* 2023; 24 (18): 13874. <https://doi.org/10.3390/ijms241813874>
49. Panche AN, Diwan AD, Chandra SR. Flavonoids: an overview. *J Nutri Sci.* 2016; 5: e47. doi: 10.1017/jns.2016.41. PMID: 28620474; PMCID: PMC5465813.
50. Passos CP, Domingues MR, Silva CM, Coimbra MA. Book Chapter -Tannins: Types, Foods Containing, and Nutrition, Nova Science Publishers. 2012.
51. Ashok PK, Upadhyaya K. Tannins are Astringent. *J Pharmacog Phytochem.* 2012; 1 (3): 45- 50.
52. Adamczyk B, Simon J, Kitunen V, Adamczyk S, Smolander

- A A Comparative Evaluation of the Antioxidant and
A. Tannins and Their Complex Interaction with Different Organic
Nitrogen Compounds and Enzymes: Old Paradigms versus Recent
Advances. *Chemistry Open*. 2017; 16; 6(5): 610-614. doi:
10.1002/open.201700113. PMID: 29046854; PMCID: PMC5641916.
53. Amarowicz R. Tannins: the new natural antioxidants? *Europ
J Lip Sci Technol*. Editorial. 2007; 109: 549–551.
54. Singh AP, Kumar S. Applications of Tannins in Industry.
Tannins - Structural Properties, Biological Properties and Current
Knowledge. *Intech Open*. 2020. Available at:
<http://dx.doi.org/10.5772/intechopen.85984>.
55. Tong Z, He W, Fan X, Guo A. Biological Function of Plant
Tannin and Its Application in Animal Health. *Front Vet Sci*. 2022; 8:
803657. doi: 10.3389/fvets.2021.803657. PMID: 35083309; PMCID:
PMC8784788.
56. Britannica T. Editors of Encyclopaedia. alkaloid.
Encyclopedia Britannica. (2025).
<https://www.britannica.com/science/alkaloid>
57. Sirin S, Dolanbay SN, Aslim B. Role of plant derived
alkaloids as antioxidant agents for neurodegenerative diseases, *Health
Sci Rev*. 2023; 6:100071.
58. Pooja SR. In-Vitro Antioxidant activity of *Ocimum
sanctum* Linn. *Int J Tech Res Appl*. 2018; 6 (3): 47-54. www.ijtra.com
59. Nasution H, Wijaya F, Ellsworth. Characterization of
antioxidant from daunkemangi (*Ocimum sanctum*) extracted using
ultrasonic bath. *J Phy Con Ser*. 2020; 1542 012016. DOI
10.1088/1742-6596/1542/1/012016
60. Saravanan R, Ramamurthy J. Evaluation of Antioxidant
Activity of *Ocimum Sanctum*-An In Vitro Study. *Int J Dent Oral Sci*.
2021; 8(11): 5001-5005.
61. Lobo V, Patil A, Phatak A, Chandra N. Free radicals,
antioxidants and functional foods: Impact on human health. *Pharmacog
Rev*. 2010; 4(8): 118-26. doi: 10.4103/0973-7847.70902. PMID:
22228951; PMCID: PMC3249911.
62. Jyoti S, Satendra S, Sushma S, Anjana T, Shashi S.
Antistressor activity of *Ocimum sanctum* (Tulsi) against experimentally
induced oxidative stress in rabbits. *Methods Find in Exp Clin
Pharmacol*. 2007; 29(6): 411-6. doi: 10.1358/mf.2007.29.6.1118135.
PMID: 17922070.