

Review Article

Amelioration of Colitis from Nature and its Immunological Implications: Current and Future Perspectives

Emmanuel O. Adediran^{1,2*}, Prabha S. Awale³, Udai P. Singh⁴

¹Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Obafemi Awolowo University, Ile Ife, Osun State, Nigeria

²Department of Chemistry, College of Arts and Sciences, Georgia State University, Atlanta, United States of America

³Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, Idaho State University, Pocatello, United States of America

⁴Department of Pharmaceutical Sciences, College of Pharmacy, The University of Tennessee Health Science Centre, Memphis, United States of America

Received: 06.06.2021; Accepted: 03.03.2022

Abstract

Inflammatory Bowel Disease (IBD) is a chronic immune-mediated inflammatory condition of the intestine. IBD is characterized by abdominal pains, diarrhea, fever, chills, cramps, and bloating and if not properly managed, it can be life-threatening. Interestingly, medicinal plants have been identified and validated to attenuate this condition due to the presence of natural products using different animal models via aryl hydrocarbon receptor and Adenosine 5'-monophosphate (AMP)-activated protein kinase (AMPK) signaling pathways. This review briefly discusses some of the natural products and medicinal plants, that can be used and developed as therapeutics for IBD treatment coupled with their immunological consequences. It recommends the need to use computational approaches to identify novel targets as well as the synthesis of structural analogs of endogenous ligands and natural products that are modulators of the identified and novel molecular targets coupled with the profiling of their biological activities and side effects.

Keywords: Colitis, Medicinal plants, Natural products, Aryl hydrocarbon receptor, Immunology

***Corresponding Author:** Emmanuel O. Adediran, Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Obafemi Awolowo University, Ile Ife, Osun State, Nigeria. Email: emmanueladediran48@gmail.com or eadediran1@gsu.edu.

Please cite this article as: Adediran EO, Awale PS, Singh UP. Amelioration of Colitis from nature and its immunological implications: Current and Future perspectives. *Herb. Med. J.* 2022;in press.

Introduction

Crohn's disease (CD) and ulcerative colitis (UC), are two major forms of Inflammatory Bowel Disease (IBD) a chronic immune-mediated inflammatory condition of the intestine. IBD, is characterized by abdominal pains, diarrhea, fever, chills, cramps, and bloating and if not properly managed, it can be life threatening (89). As a matter of fact, it has affected millions of people and its prevalence is on the

increase everyday (1). IBD has been considered to be an autoimmune disease; a situation where the immune system dysregulates in certain part of the gastrointestinal tract and induce inflammation. (2,3). Ulcerative colitis may begin in the rectum and extend to the rest of the colon while Crohn disease can occur anywhere in the gastro intestinal tract. However, other organ systems may be involved. Although colitis is of inflammatory origin, there may be interplay of bacterial infections such as *E coli*, *Salmonella spp*,













Camphylobacter spp, parasites and viruses which will invariably demand for an antibiotic therapy. Surprisingly, genetic and environmental influences are also involved in the pathogenesis of colitis but its etiology is not fully understood (3). Interestingly, because of the diversity of nature, medicinal plants have been a platform for the treatment of various inflammatory diseases in

different cultures of the world. As a matter of fact, medicinal plants have been scientifically proven to possess ranges of beneficial pharmacological and therapeutic purposes ranging from antiviral, antioxidant, anti-inflammatory, antiinflammatory, antihypertensive, immunostimulating, cytotoxic and antibacterial properties (4, 5). Furthermore, the highlighted

Table 1: Pharmacological and biological activities of selected medicinal plants and reported natural products.

Medicinal Plants	Morphological Parts	Compounds Isolated	Reported Biological Activities	References
<i>Anacardium occidentales</i>	leaves, cashew fruit, cashew apple juice	Agathisflavone, Quercetin-3-O-rhamnoside, Anacardic acids, 2-methylcardols	Antioxidant, antibacterial, cytotoxic	(6-8)
<i>Ixora coccinea</i>	Flowers, leaves, roots	Ixoroid Ursolic acid Ixoratannin A-2 Procyanidin A2 Cinnamtannin B1	Antioxidant Antimicrobial Anti-genotoxic Antinociceptive Catarrhal bronchitis Antileishmanial	(9-11)
<i>Terminalia arjuna</i>	Bark	Procyanidine, Arjunolic acid	Molluscicidal Anti acetylcholinesterase	(12)
<i>Vernonia amygdalna</i>	Flowers, leaves	Tricosane Vernolide Isorhamnetin Luteolin Vernolide Vernodalol	Antioxidant Neuroprotective Anti-alzheimer Anti-microbial Anti-lipid	(13-15)
<i>Camella sinensis</i>	leaves, flowers	Teadenol A Teadenol B Caffeine, myricetin-rhamnoglucoside teasperol teasperin theaflavins thearubigins	Antioxidant, Antibacterial Cytotoxic Anticaries Antiparkinson Cardiovascular Anti-HIV	(16-20)
<i>Chochorus olitorius</i>	Leaves	Caffeic acid Chlorogenic acid Isorhamnetin	Antioxidant Hypoglycemic Hypolipidaemic Anti-hypertension Immunomodulatory	(21-24)
<i>Mangifera indica</i>	Stem bark, leaves, Seed	Tricuspid Mangiferin Tetragalloylglucose Ellargic acid Iriflophenone glucoside	Anti-fungi Analgesic Anti-inflammatory Antioxidant Immunomodulatory antiplatelet	(25-28)
<i>Curcuma longa</i>	Rhizome	Curcumin	Anti-inflammatory	(29, 30)
<i>Azadirachta indica</i>	Stem bark, leaves	Neemazal Azadirachtin A Azadirachtin B Nimbin Salannin	Antimicrobial Antimalaria	(85, 86)
<i>Zingiber officinales</i>	Rhizome	Terpens 4-gingerol 6-dehydrogingerdione	Antioxidant Antiinflammatory Anticoagulant Antimicrobial	(87, 88)

Figure 1. Classification of flavonoids with specific examples.

Flavonoid Classes	General Structures	Specific Examples
FLAVONE		 APIGENIN
FLAVONOL		 QUERCETIN
FLAVANONE		 HESPERETIN
ISOFLAVONE		 GENISTEIN
ANTHOCYANIDIN		 CYANIDIN
CHALCONE		 PHLORETIN

This figure is modified (31)

pharmacological and biological activities are mediated by the presence of one or more natural products and metabolites in the plant ranging from flavonoids, saponins, tannins, alkaloids, polyphenols, anthraquinones. This claims have repeated been

validated by phytochemical screenings, bioassay guided fractionations and the use of animal models (6). In this review we will briefly discuss on some of the natural products and medicinal plant that can be use and develop as therapeutics for IBD treatment




Figure 2. Immunological response mediated by the activation of Aryl hydrocarbon receptor by a dietary flavonoid.

coupled with their immunological consequences.

Flavonoids and its diversity

One of the most common alternative medicine compound groups comes from Flavonoids. Flavonoids are a group of polyphenolic compounds, diverse in chemical structure and characteristics, found ubiquitously in many plants. Therefore, flavonoids are part of the human diet in many countries around the world. The structural requirements for the antioxidant and free radical scavenging functions of flavonoids include a hydroxyl group in carbon position three, a double bond between carbon positions two and three, a carbonyl group in carbon position four, and polyhydroxylation of the A and B aromatic rings (32). Flavonoids also widely occur in Fabaceae, Polygonaceae and Rosaceae families.

Flavonoids have been identified to ameliorate colitis. Specifically, Extracts containing eupatilin and quercetin-3- β -D-glucuronopyranoside ameliorated the inflammatory response and colonic injury in acute colitis by decreasing oxidative stress and neutrophil activation (90).

There are different classes of flavonoids ranging from flavones, flavanones, flavonols, isoflavones, biflavonols and each of the chemical compounds have a characteristic modulatory effect of the aryl hydrocarbon receptor; a target that is indicated in the inflammatory pathway of the IBD as well as in many autoimmune diseases. Some of the compounds are proven to be inhibitory and some are stimulatory (32). Hence, there is a need to identify medicinal plants that are more richer in the flavonoids with stimulatory effect which can invariably be of help in the formulation of herbal medicines.

Aryl Hydrocarbon Receptor

Aryl hydrocarbon receptor is a transcription factor embedded in the cytosol with functioning domains

and binding sites (33, 34). Reports have shown that there is high expression of it in the liver, kidney, placenta, lungs, thymus (35). The AhR and its nuclear translocator (ARNT) belong to the bHLH-PAS superfamily of transcription factors, the acronym for basic Helix-Loop-Helix and Periodic (PER)-ARNT Single minded (SIM) (36). AhR is capable of exhibiting a conformational changes and its ability of translocation to the nucleus to bind to its dimerization partner; nuclear translocator forms a complexes once activated by an agonist (37-39). The formation of complexes initiates the transcription of genes and other molecular events.

Aryl hydrocarbon receptor has been an evolving pharmacological target in the management in cancer, inflammation, autoimmune diseases and ageing (40, 41). It has plays a role in the induction of cytochrome P450 enzymes and other metabolizing enzymes such as cytosolic phospholipase and cyclooxygenase enzymes. Immunological, carcinogenic effects, inflammatory responses, endocrine disruptions has been linked to Aryl hydrocarbon receptor mediated activity (42). It has also been implicated in key physiological functions such as cell cycle regulations, resilience against stress, homeostasis, immunity and regulation of neural functions in vertebrate and invertebrates (43, 44). The functional diversities of this receptor has made it to be tagged a "promiscuous receptor" of chemical entities (45). However, the activity mediated by this receptor is ligand dependent and calls for a profiling of natural products in this regard. For example, when comparing the activity of apigenin, a flavone with its analogue genistein; an isoflavone there is a loss of agonistic effect. A modification from position 2 to position appears to be the reason for the loss of the agonist effect (32). These non-general perspective was further strengthened when the interaction of 2,3,7,8-tetrachlorodibenzo-p-

dioxin with aryl hydrocarbon receptor resulted into reduced experimental autoimmune encephalomyelitis (EAE) by promoting the development of Foxp3+ Treg cells, whereas 6-formylindolo(3,2-b) carbazole enhances EAE by inducing the differentiation of IL-17-producing T cells (46-48).

Immunological Dysregulations

In a bid to have an understanding of the immunological consequences associated with inflammation in IBD, it involves the use of different animal models. The approach has been to specifically delete the Aryl hydrocarbon receptor in T cells or Macrophages in mice and also the use of dextran sulphate sodium(DSS) or trinitrobenzenesulfonic acid (TNBS), to induce colitis in mice (49,50,51). Aryl hydrocarbon receptor is expressed in normal T cells, dendritic cells, macrophages and cells in the gut.

T Cells and Functions

T cells are vital player in the regulation of adaptive immunity, immune responses and tissue inflammation. Experimental and clinical evidence have also shown that chemokines and cytokines plays a crucial role in the regulation of colitis by promoting the migration of leucocytes to the site of inflammation leading to tissue disruption and damage (52). Data have also shown that there is a significant increase in the circulating level of specific chemokines and cytokines of colitis patients when compared to normal healthy subjects (52). The specific inflammatory cytokines are IL-16, IFN- γ , IL-1 β and TNF- α (52). The inhibition of the activities of the inflammatory mediators can be explored to either ameliorate or reverse colitis symptoms.

Most major T cell types are expressed in the thymus, although extrathymic generation of some T cell subsets has been described (53, 54). T cells are classified into the major TCR- α β and minor γ δ T cell groups. $\alpha\beta$ T cells are highly heterogeneous and grouped into CD4+ conventional T cells, CD8+

conventional T cells, NKT cells, and other innate TCR α - expressing T cells such as mucosal-associated invariant T (MAIT) cells (55,56,57,58). CD4+ conventional T cells are further divided into FoxP3+ regulatory and FoxP3- T cells (59,60). FoxP3- CD4+ T cells include various effector and regulatory T cells based on their cytokine phenotype (IFN γ , IL-17, IL-22, IL-4, IL-9, IL-10, IL-35, and/or LAP-TGF β 1) (60,61). These T helper cells include IFN γ + Th1 cells, IL-17/IL-22+ Th17 cells, IL-4+ Th2 cells, IL-9+ Th9 cells, IL-21+ T-FH cells, and IL-10/IL-35/TGF β 1+ Tregs (61,62). All of these T helper cell subsets are generated mainly in the periphery from naïve T cells made in the thymus.

Driven by the conviction to dig into nature, *Pycnoporus sanguineus* (L) Murril is a saprotrophic fungus. Recently, polysaccharides from the plant have been reported to ameliorate colitis which manifested by the lowered disease activity index (81). The mechanism of action was also documented to be due to inhibition of helper T cells mediated immune response by decreasing the proportions of Th cells (including Th2 cells, Th 17 cells and regulatory T cells) with an accompanied reduction in myeloperoxidase (MPO) activities and release of several interleukins and chemokines within the colon.

Activation of aryl hydrocarbon receptor, a transcription factor by flavonoids plays a considerable role in the regulatory T cell (Treg) induction (63). The induction of this T cells regulate and suppress allergies and autoimmune disease; Crohn disease as a typical example

The activation of Aryl hydrocarbon receptor by flavonoids and its ability to ameliorate inflammatory conditions is also believed to be due to its free radicals scavenging effects, inhibition of oxidative stress and attenuation of Reactive Oxygen Species(ROS) (64, 65).

Table 2: Biological and Immunological Implications of Selected Natural Products.

Natural Products	Targets	Modulatory Effects	biological and immunological implications	References
Beberine	AMPK	Activation	1. Inhibition of oxidative phosphorylation 2. BBR suppressed inflammatory cytokines of LP CD4+ T cells from colitis SciD mice in vitro. 3 BBR increased the AMPK activity and regulated the IFN- γ and IL-17A secretion from colitis LP CD4+ T cells.	(68, 75)
Narigenin	Aryl hydrocarbon Receptor	Activation	Induction of regulatory T Cells	(63)
Resveratrol	Aryl hydrocarbon receptor, AMPK	Activation	Reduction of COX-2 expression, neutrophil mobilization Decrease mitochondrion fatty acid oxidation Decrease IL 6 expression and TNF-alpha release Activation of myeloid derived suppressor cells	(49, 75)
Quercetin	AMPK	Activation	Amelioration of IgE-mediated allergic intestinal inflammation	(75, 76)
Sinapic Acid	Antioxidant enzymes	Activation	Enhancing the activities of superoxide dismutase, glutathione peroxidase. Inhibition of Myeloperoxidase activities	(77)
Myricetin & Myricetin-3-O-B-D-lactose sodium salt	Intestinal microbiota	Normalization	Normalization of proportion of firmicutes and actinobacteria. Inhibition of pathogenic microorganisms such as Ruminococcus and Parabacteroides. Increase in probiotics such as akkermansia	(78)
Curcumin	Proinflammatory cytokines	Inhibition	Inhibition of IFN and TNF alpha Inhibition of COX-2 Obstruction of p38 MAPK signaling Reduction in the Infiltration of leucocytes	(29, 30, 75)
Epigallocatechin-3-gallate	Proinflammatory cytokines	Inhibition	Increase in antioxidant enzymes (SOD and GPO) activities. Inhibition of the production of proinflammatory cytokines	(75-80)

It is interesting that another new target in the pathogenesis of chronic colitis has been identified and reported (66-68). AMPK is an enzyme that plays a role different fundamental cellular processes, including the cell proliferation, survival and metabolism and in cellular energy homeostasis (69-71). Energy metabolism changes in immune cells have been identified to be crucial for immune cells regulation. Recently, it was reported that Berberine, a traditional Chinese herbal medicine extracted from *Phellodendron* bark and *Coptis japonica* ameliorate chronic colitis through the activation of AMPK, inhibition of oxidative phosphorylation and suppression of the exaggeration of IFN- γ - and IL-17A-producing LP CD4+ T cells (68). Novel animal

model (T cell transfer colitis model) which involves the transfer of naïve (CD4+CD45RB^{high}) T cells into congenic immunodeficiency mice to induce CD4+ T cell-specific colitis were used to validate the claim (68, 72). Infact, it reflects the chronic inflammatory process. Although, several reports have used chemical agents like DSS to induce colitis (73, 74), it can only be used for acute colitis models.

Biological and Immunological Implications of Selected Natural Products

Carbon Monoxide (CO); an endogenously produced signaling molecules has been identified to be a promising therapeutic agent against inflammation and some form of cellular stress related diseases (83, 84). Although it is a pollutant produced by the incomplete combustion of carbon, it is a “glorified” pollutant.

Reports have shown its ability to ameliorate colitis and effort are ongoing to obtain its pharmacokinetic and clinical data and its delivery form optimization. CO was shown to downregulate pro-inflammatory cytokines TNF- α , IL-1 β (82, 84).

Conclusion

Although, Aryl hydrocarbon receptor, AMPK are vital molecular targets in the pathogenesis of colitis, Computational approaches can be leveraged on to identify additional novel promising targets, multiple activators of receptor sites which can be characterized and exploited in drug design and discovery. A synergistic effect of chemical compounds that are of therapeutic value will be of help in this regard.

Furthermore, synthesis of structural analogues of endogenous ligands and natural products that are modulators of the identified molecular targets can be launched coupled with the profiling of their biological activities and side effects.

This report has shown the beauty of nature, target diversity and the abundant of natural products. Medicinal plants can be repurpose for this indication; colitis and some other known and evolving autoimmune diseases.

The medicinal value of dietary supplements, fruits, herbal medicines will complement the orthodox drugs in the management of colitis and other inflammatory disease to improve the quality of life of the patients and if need be, there can be a switch to nature if the side effects of orthodox drugs are pronounced.

Acknowledgments

The authors acknowledge His-kai Wang for the diagrammatic illustration of the immunological implications of Narigenin.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

1. Zhou M., He J, Shen Y, Zhang C, Wang J, Chen Y. New

- frontiers in genetics, gut microbiota, and immunity: A rosetta stone for the pathogenesis of inflammatory bowel disease. *Biomed. Res. Int.* 2017;1–17. (CrossRef) (PubMed)
2. Podolsky DK. The current future understanding of inflammatory bowel disease. *Best Pract Res Clin Gastroenterol.* 2002;16:933-43.
3. Bouma G, Strober W. The immunological and genetic basis of inflammatory bowel disease. *Nature Reviews Immunology.* 2003;3:521-33.
4. Vazirian M, Nabavi SM, Jafari S, Manayi A. Natural activators of adenosine 5'-monophosphate (AMP)-activated protein kinase (AMPK) and their pharmacological activities. *Food and Chemical Toxicology.* 2018.
5. Thea M, Manrico M, Matheo A R, Emilio J. Recent Advances on the anti-inflammatory and anti-oxidant properties of Red Grape Polyphenols: in vitro and in vivo studies. *MDPI.* 2020;9(1):35.
6. Ajileye EM, Obuotor EO, Akinkunmi MA. Aderogba Isolation and characterization of antioxidant and antimicrobial compounds from *Anacardium occidentale* L.(Anacardaceae) leaf extract. *Journal of King Saud University-Science:* 2015;27(3):244-52.
7. Isaokubo Masamitsu ochi, Paulo C Vieira, Sakae Komatsu. Antitumor agents from the cashew apple juice. *Journal of Agricultural and food chemistry.* 1993;41(6):1012-5.
8. Isao K, Ikuyo K, Yoshihiro Y. Tyrosine inhibitors from *Anacardium occidentale* fruit. *Journal of Natural products.* 1994;57(4), 545-51.
9. Thomas OI, Abiodun OO, Abiola OS, Efere M, et al. Doubly linked A-type proanthocyanidin trimer and other constituents of *ixora coccinea* leaves and their antioxidant and antibacterial properties. *Journal of phytochemistry.* 2010;71(17-18):2092-8.
10. CY Ragasa, F Tiu, JA Rideont. New cycloartenol esters from *ixora coccinea*. *Natural product research.* Taylor and Francis. 2004.
11. Haji M, Muhammed Q, Ambreen I, Versiani MA, Tahiri IA, et al. Antioxidant antimicrobial activities of *ixora coccinea* root and quantification of phenolic compounds using HPLC. *South African Journal of Botany.* 2020;135:71-9.
12. Neelam S, Dinesh K, Singh V, Kumar S. Inhibition kinetics of acetylcholinesterase and phosphatases by the active constituents of *Terminalia arjuna* and *Tamarindus indica* in the cerebral ganglion of *Lymnaea acuminata*. *Journal of Pharmacognosy.* 2017;9(2).
13. Ganiyu O, Bukola Ch, Mayowa BA, Idowu S, Opeyemi B. Characterization and neuroprotective properties of alkaloids extract of *vernonia amydalyna* Delile in Experimental models of Alzheimer disease. *Drug and Chemical Toxicology.* 2020;1-10.
14. Abere H, Yadessa M. Antibacterial and antioxidant compounds from the flower extracts of *vernonia amygdalina*. *Advances in Pharmacological sciences.* 2018.
15. P Erasto DS, Grierson AJ, Afolayan. Antioxidant constituents in *vernonia amydalyna* leaves. *Pharmaceutical biology.* 2007;45(3).195-9.
16. Ali I, Masood SB, et al. Inhibitory effect of black tea (*Camellia sinensis*) theaflavins and thearubigins against HCT 116 colon cancer cells and HCT 460 lung cancer cells. *Journal of food biochemistry.* 2019;43(5):212822.
17. Ayaka K, Arisa S, et al. New phenolic compounds from *Camellia sinensis* L. fermented leaves. *Journal of natural medicines.* 2013;67(3):652-6.
18. Taylor PW, Hamilton-Miller JM, Stapleton PD. Antimicrobial properties of green tea catechins. *Food Sci Technol Bull.* 2005;2:71-81 (pubmed:19844590).
19. Bertoldi M, Gonsalvi M, Voltattorni CB. Green tea polyphenols: Novel irreversible inhibitors of dopa decarboxylase. *Biochem. Biophys. Res. Commun.* 2001;284:90-3.
20. Xin W, Quan L, Hongho Zh, Hongqing W. Flavanols from the *Camellia sinensis* var. *assamica* and their hypoglycemic and hypolipidemic activities. *Acta pharmaceutica sinica B.* 2017;7(3):342-6.

21. Augustine I, Airaodion, Emmanuel O, Ogbonnaya E, Uloaku O, Olaide A, Olukule A. Ameliorative efficacy of phytochemical content of corchorus olitorius leaves against acute ethanol-induced oxidative stress in wistar rats. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2019;1-10.
22. Augustine I, Emmanuel O, Ogbonnaya E, Uloaku O, Olaide A, Olukule A. Corchorus olitorius leaves on hypoglycemic and hypolipidaemic activities in albino rats. *Asian Plant Research Journal*. 2019;1-13.
23. Ganiyu O, Adedayo O, Ayodele J, Thomas H, Jamiyu A, Uine S. Inhibitory effect of polyphenol-rich extracts of jute leaf (corchorus olitorius) on key enzyme linked to type 2 diabetes (alpha-amylase and alpha-glucosidase) and hypertension. *Journal of functional foods*. 2012;4(2):450-8.
24. Ho-Young P, Mi-jinoh Y, Inwook Ch. Immunomodulatory activities of corchorus olitorius leaf extract: Beneficial effects in macrophage and NK cell activation immunosuppressed mice. *Journal of Functional foods*. 2018;46:220-6.
25. Qudsia K, Ishtiaq H, Siddiqui HL, Javaid A. Antifungal activity of flavonoids isolated from mango (*Mangifera indica* L) leaves. *Natural product Research*. 2010;24(20):1907-14.
26. Gabino G, Deyarina G, Carla D. Analgesic and anti-inflammatory effects of mangifera indica L. extract (vimang). *Phytotherapy Research: An international journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*. 2001;15(1):18-21.
27. Rodeiro I, Cancino L, Gonzalez J E, Morffi J, et al. Evaluation of the genotoxic potential of *Mangifera indica* L extract, a new natural product with antioxidant activity. *Food and Chemical Toxicology*. 2006;44(10):1707-13.
28. Maria Elena A, Palomo M, Rodriguez L, Fuentes E. Antiplatelet activity of natural bioactive extracts from mango and its by products. *Antioxidants*. 2019;8(11):517.
29. Yasuyuki D, Akira A, Osamu I, Yuhki Y, Shigeki B, Yoshio A, et al. Curcumin prevents the development of dextran sulfate sodium (DSS)-induced experimental colitis. *Digestive diseases and sciences*. 2007;52(11):2993-8.
30. Karthikeyan A, Young KN, Moniruzzaman M, Beyene AM, Do K, Kalaiselvi S, et al. Curcumin and Its Modified Formulations on Inflammatory Bowel Disease (IBD): The Story So Far and Future Outlook. *Pharmaceutics*. 2021;13:484.
31. Goya-Jorge E, Jorge Rodríguez ME, Veitía MS, Giner RM. Plant Occurring Flavonoids as Modulators of the Aryl Hydrocarbon Receptor. *Molecules*. 2021;26:2315.
32. Harborne JB, Baxter H. *Handbook of Natural Flavonoids*. Chichester: Wiley. 1999.
33. Perdew GH. Association of the Ah receptor with the 90-kDa heat shock protein. *J. Biol. Chem*. 1988; 263:13802.
34. Sogawa K, Fujii-Kuriyama Y. Ah receptor, a novel ligand-activated transcription factor. *J. Biochem*. 1997;122:1075.
35. Carver LA, Hogenesch JB, Bradfield CA. Tissue Specific Expression of the Rat Ah-Receptor and ARNT mRNAs. *Nucleic Acids Res*. 1994;22:3038-3044.
36. Hogenesch JB, Chan WK, Jackiw VH, Brown RC, Gu YZ, Pray-Grant M, et al. Characterization of a Subset of the Basic-Helix-Loop-Helix-PAS Superfamily That Interacts with Components of the Dioxin Signaling Pathway. *J. Biol. Chem*. 1997;272:8581-93.
37. Burbach KM, Poland A, Bradfield C. A. Cloning of the Ah-receptor cDNA reveals a distinctive ligand-activated transcription factor. *Proc. Natl Acad. Sci*. 1992; 89:8185.
38. Larigot L, Juricek L, Dairou J, Coumoul X. AhR Signaling Pathways and Regulatory Functions. *Biochim*. 2018;7:1-9.
39. Schulte KW, Green E, Wilz A, Platten M, Daumke O. Structural Basis for Aryl Hydrocarbon Receptor-Mediated Gene Activation. *Structure*. 2017;25:1025-33.
40. Lamas B, Natividad JM, Sokol H. Aryl Hydrocarbon Receptor and Intestinal Immunity. *Mucosal Immunol*. 2018;11:1024-38.
41. Gao J, Xu K, Liu H, Liu G, Bai M, Peng C, et al. Impact of the Gut Microbiota on Intestinal Immunity Mediated by Tryptophan Metabolism. *Front. Cell. Infect. Microbiol*. 2018;8:1-22.
42. Denison MS, Soshilov AA, He G, Degroot DE, Zhao B. Exactly the Same but Different: Promiscuity and Diversity in the Molecular Mechanisms of Action of the Aryl Hydrocarbon (Dioxin) Receptor. *Toxicol. Sci*. 2011;124:1-22.
43. Mescher M, Haarmann-Stemmann T. Modulation of CYP1A1 Metabolism: From Adverse Health Effects to Chemoprevention and Therapeutic Options. *Pharmacol. Ther*. 2018;187:71-87.
44. Juricek L, Coumoul X. The Aryl Hydrocarbon Receptor and the Nervous System. *Int. J. Mol. Sci*. 2018;19:2504.
45. Stejskalova L, Dvorak Z, Pavek P. Endogenous and Exogenous Ligands of Aryl Hydrocarbon Receptor: Current State of Art. *Curr. Drug Metab*. 2011;12:198-212.
46. Veldhoen M, Hirota K, Westendorf AM. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. *Nature*. 2008;453:106.
47. Quintana FJ, Basso AS, Iglesias AH. Control of T(reg) and T(H)17 cell differentiation by the aryl hydrocarbon receptor. *Nature*. 2008;453:65.
48. Nam Trung Nguyen, Hamza Hanieh, Taisuke Nakahama and Tadimitsu Kishimoto. The roles of aryl hydrocarbon receptor in immune responses. *International Immunology*. 2013;25;6:335-43.
49. Udai PS, Narendra PS, Balwan S, Lorne JH, Robert L. Resveratrol (Trans-3,5,4-trihydroxystilbene) Induces Silent Mating Type Information Regulation-1 and Down-Regulates Nuclear Transcription Factor B Activation to Abrogate Dextran Sulfate Sodium-Induced Colitis. *Journal of Pharmacology and experimental therapeutics*. 2009;332:829-39.
50. Singh NP, Singh UP, Singh B, Price RL, Nagarkatti M, et al. Activation of Aryl Hydrocarbon Receptor (AhR) Leads to Reciprocal Epigenetic Regulation of FoxP3 and IL-17 Expression and Amelioration of Experimental Colitis. *PLoS ONE*. 2011; 6(8): e23522. doi:10.1371/journal.pone.0023522.
51. Egger B, Bajaj-Elliott M, MacDonald TT, Inglin R, Eysselein VE, et al. Characterisation of acute murine dextran sodium sulphate colitis: cytokine profile and dose dependency. *Digestion*. 2000; 2:240-8.
52. Udai PS, Narendra PS, Angela M, Robert L. Chemokines and cytokines level in inflammatory bowel disease patients. *Elsevier Journal. Cytokines*. 2015;77:44-49.
53. Heinonen KM, Perreault C. Development and functional properties of thymic and extrathymic T lymphocytes. *Crit. Rev. Immunol*. 2008;28:441-66.
54. Bhandoola A, Von BH, Petrie HT, Zuniga-Pflucker JC. Commitment and developmental potential of extrathymic and intrathymic T cell precursors: plenty to choose from. *Immunity* 2014;26:678-89.
55. Gapin L, Check M. *J Immunol*. 2014;192: 4475-80.
56. Rossjohn J, Pellicci DG, Patel O, Gapin L, Godfrey DI. Recognition of CD1d-restricted antigens by natural killer T cells. *Nat. Rev. Immunol*. 2012;12:845-57.
57. Wan YY. Multi-tasking of helper T cells. *Immunology*. 2010;130:166-71.
58. Li P, Spolski R, Liao W, Leonard WJ. Complex interactions of transcription factors in mediating cytokine biology in T cells. *Immunol. Rev*. 2014;261:141-56.
59. Gratz IK, Campbell DJ. Organ-specific and memory treg cells: specificity, development, function, and maintenance. *Immunol*. 2014;5:333.
60. Liston A, Gray DH. Homeostatic control of regulatory T cell diversity. *Nat. Rev. Immunol*. 2014;14:154-65.
61. Liu X, Nurieva RI, Dong C. Transcriptional regulation of

- follicular T-helper (T_{fh}) cells. *Immunol. Rev.* 2013;252:139-45.
62. Tripathi SK, Lahesmaa R. Transcriptional and epigenetic regulation of T-helper lineage specification. *Immunol. Rev.* 2014;261:62-83.
63. His-kai W, Chen-Hao Y, Taku I, Hideosatsu MSh, Mamoru T. Dietary flavonoids Naringenin induces Regulatory T cells via aryl hydrocarbon receptor mediate pathway. *Journal of Agricultural and Food Chemistry.* 2012;60(9):2171-8.
64. Abotaleb M, Samuel SM, Varghese E, Varghese S, Kubatka P, Liskova A, et al. Flavonoids in Cancer and Apoptosis. *Cancers.* 2019;11:28.
65. van der Heiden E, Bechoux N, Muller M, Sergent T, Schneider YJ, Larondelle Y, et al. Food Flavonoid Aryl Hydrocarbon Receptor-Mediated Agonistic/Antagonistic/Synergic Activities in Human and Rat Reporter Gene Assays. *Anal. Chim. Acta.* 2009;637:337-45.
66. Rao E. AMPK-dependent and independent effects of AICAR and compound C on T-cell responses. *Oncotarget.* 2016;7:33783-95.
67. Blagih J. The energy sensor AMPK regulates T cell metabolic adaptation and effector responses in vivo. *Immunity.* 2015;42:41-54.
68. Masahiro T, Akinobu T, Sakiko H, Takuya A, Yasuyuki Sh, Hiroshi M, et al. Berberine improved experimental chronic colitis by regulating interferon- γ - and IL-17A-producing lamina propria CD4⁺ t cells through AMPK activation.scientific report;nature research. 2019.
69. Liu Y. Berberine inhibits macrophage M1 polarization via AKT1/SOCS1/NF-kappaB signaling pathway to protect against DSS-induced colitis. *International immunopharmacology* 2018;57:121-31.
70. Ganeshan K, Chawla A. Metabolic regulation of immune responses. *Annual review of immunology.* 2014; 32:609-34.
71. Loftus RM, Finlay DK. Immunometabolism: Cellular Metabolism Turns Immune Regulator. *The Journal of biological chemistry.* 2016;291:1-10.
72. Gubser PM. Rapid effector function of memory CD8⁺ T cells requires an immediate-early glycolytic switch. *Nature immunology.* 2013;14:1064-72.
73. Yan F. Berberine promotes recovery of colitis and inhibits inflammatory responses in colonic macrophages and epithelial cells in DSS-treated mice. *American journal of physiology. Gastrointestinal and liver physiology.* 2012;302:504-14.
74. Yu XT. Berberine attenuates mucosal lesions and inflammation in dextran sodium sulfate-induced colitis in mice. *PloS one.* 2018;13.
75. Vazirian M, Nabavi SM, Jafari S, Manayi A., Natural activators of adenosine 5'-monophosphate (AMP)-activated protein kinase (AMPK) and their pharmacological activities, *Food and Chemical Toxicology.* 2018.
76. Yuanyang D, Qihang H, Jiaqu L, Patricia G. Wolf, Hamed Ayansola and Bingku Zhang. Quercetin alleviate intestinal oxidative Damage induced by hydrogen peroxide via modulation of GSH;invitro screening and in vivo evaluation in a colitis model.*American Chemical Society.* 2020;5(14):8334-48.
77. Bo Qian. Ameliorative effect of sinapic acid on Dextran Sodium Sulfate(DSS) induced ulcerative colitis in Kunming KM mice.*Oxid Med Cell Longev.* 2020.
78. Miao RR, Zhan S, Hu X T,Yuan W M. .Myricetin and M10,a myricetin-3-B-D lactose sodium salt,modifying composition of gut microbiota in mice with ulcerative colitis.*Toxicology letters.* 2021;346:7-15.
79. Markus B, SabineK. Westphal W, Domschke TK. Green tea polyphenol epigallocatechin-3-gallate shows therapeutic antioxidative effects in a murine model of colitis.*Journal of Crohns and Colitis.* 2012;6(2):226-35.
80. Brittany A. Payan K, Vanarsa H. Epigallocatechin-3-gallate ameliorates dextran sulfate sodium(DSS)-induced inflammatory bowel disease. *J.Immunol.* 2018;1 supplement:54.12.
81. Muxia Li,Ting Luo,Young Huang etal. Polysaccharide from *Pycnoporus Sanguineus* ameliorate dextran Sulfate Sodium induced colitis via helper T cells repertoire modulation and autophagy suppression. 2020;34(10):2649-64.
82. Marks GS, Brien JF, Nakatsu K, et al. Does carbon monoxide have a physiological function? (J). *Trends Pharmacol Sci,* 1991, 12(5): 185-188.
83. YANG Xiao-Xiao, KE Bo-Wen, LU Wen, WANG Bing-He. CO as a therapeutic agent: discovery and delivery forms. *Chinese Journal of Natural Medicines.* 2020;18(4):284-95.
84. Takagi T, Uchiyama K, Naito Y.The therapeutic potential of Carbon Monoxide for inflammatory bowel disease. *Digestion.* 2015;91:13-18.
85. Vidhu Sankar Babu, S Narasimhan,GM Nair.Bioproduction of Azadirachtin-A,Nimbin and salannin in callus and cell suspension cultures of neem (*Azadirachta indica* A. Juss).*Current science.* 2006;91(1):22-4.
86. E Coventry,Eunice Jessie Allan.Microbiological and Chemical analysis of neem (*Azadirchta indica*) extracts:new data on antimicrobial activity. *Phytoparasitica.* 2001;29(5):441-50.
87. Wail E, Emad M. Antibacterial activity of ginger(*Zingiber officinale* Rosc.) rhizome: a mini review. *International journal of pharmacognosy and chinese medicine.* 2018;2(4):1-8.
88. Rahuman A, Geetha G, Venkatesan P, Kannappan G, Bagavan A. Mosquito larvicidal activity of isolated compounds from the rhizome of *Zingiber officinale*.*Phytotherapy Research.* 2008;22(8):1035-9.
89. Singh BP, Marshall JL, He AR. Workup and Management of Immune-Mediated Colitis in Patients Treated with Immune Checkpoint Inhibitors. *The oncologist.* 2020;25(3):197-202.
90. Minjae Joo,Han Sang Kim,Tae Hoon Kwon, Alisha Palikhe, Tin Sandar Zaw, Ji Hoon Jeong, and Uy Dong Sohn. Anti-inflammatory Effects of Flavonoids on TNBS-induced Colitis of Rats. *Korean J Physiol Pharmacol.* 2015;19(1):43-50.

© Emmanuel O. Adediran, Prabha S. Awale, Udai P. Singh, Originally published in the *Herbal Medicines Journal* (<http://www.hmj.lums.ac.ir>), 17.05.2022. This article is an open access article under the terms of Creative Commons Attribution License, (<https://creativecommons.org/licenses/by/4.0/>), the license permits unlimited use, distribution, and reproduction in any medium, provided the original work is properly cited in the *Herbal Medicines Journal*. The complete bibliographic information, a link to the original publication on <http://www.hmj.lums.ac.ir/>, as well as this copyright and license information must be included